

Snail Bullhead

Ameiurus brunneus

Contributor (2005): Jason Bettinger (SCDNR)

Reviewed and Edited (2013): Mark Scott, Andrew R. Gelder, and M. Troy Cribb [SCDNR]



DESCRIPTION

Taxonomy and Basic Description

The Snail Bullhead is a member of the family Ictaluridae. The ictalurids, with approximately 46 species, represent the largest family of fishes indigenous to only North America (Jenkins and Burkhead 1994). The genus *Ameiurus* contains the moderately-sized (reaching only a few pounds) catfish collectively known as the “bullheads.” Snail Bullheads have a large flat head, a decurved snout profile, uniformly colored maxillary barbel, short rounded anal fin, emarginate tail and a large dusky blotch at the base of the dorsal fin. Snail Bullheads vary greatly in color and pattern, but in general they have a brown or yellow-green back, mottled sides of the same colors, and a white belly. In their natural habitats (rivers and streams), Snail Bullheads typically reach maximum adult sizes of 294 mm (11.4 in.) (Rohde et al. 1994); however, in disturbed habitat such as impounded river reaches and reservoirs, they reach substantially greater sizes (443 mm or 17.6 in.) (SCDNR unpublished data).

Status

The Snail Bullhead is apparently secure (G4) over its range, and is not listed (SNR) in South Carolina (NatureServe 2013). However, in a recent assessment of southeastern fishes, this species was considered vulnerable to imperilment (V) (Warren et al. 2000; Jelks et al. 2008). The Snail Bullhead is considered at least vulnerable to imperilment in 4 of the 6 states where it occurs. It is considered imperiled (S2) in Virginia, vulnerable (S3) to imperilment in Alabama and Florida, and apparently secure (S4) in Georgia and North Carolina (NatureServe 2013).

POPULATION SIZE AND DISTRIBUTION

Snail Bullheads occur along the Atlantic Slope from the Dan River system in Virginia and North Carolina south to the Altamaha River system in Georgia and the Saint Johns River drainage in Florida (NatureServe 2004). They also occur in Gulf Slope drainages of the Apalachicola River system in Georgia, Alabama, and Florida and in the upper Coosa River system in northern Georgia (Page and Burr 1991).

Snail Bullheads appear to be relatively stable in South Carolina streams; however, their fate in the larger rivers is uncertain due to the presence of the nonnative Flathead Catfish. They are common to abundant in the upper Santee drainage, but decreasing trends have been noted in many coastal rivers like the Edisto River, following the introduction of Flathead Catfish (C.

Thomason, SCDNR, pers. comm.). The largest populations of Snail Bullhead are probably located in the Broad River system, a system where the Flathead Catfish has yet to be introduced. Based on South Carolina Stream Assessment (2006-2011) data, the mean statewide density estimate for Snail Bullhead in wadeable streams was 0.02 (95% confidence interval: 0.01–0.03) per 100 m².

HABITAT OR NATURAL COMMUNITY REQUIREMENTS

The Snail Bullhead is found in rocky riffles, shoals, runs, and pools in streams and rivers. (Jenkins and Burkhead 1994; Rohde et al. 1994; NatureServe 2013).

CHALLENGES

The Snail Bullhead is adversely affected by the same myriad of challenges plaguing all aquatic species such as sedimentation, hydrologic modification, impoundments, non-point source pollution, and development. Additionally, this species is also threatened by the introduction of non-native ictalurids like the Flathead Catfish and the Blue Catfish; both of these non-native species displace and prey on the smaller bullheads. Flathead Catfish have been shown to prey on bullhead species and greatly reduce their numbers (Guire et al. 1984; Ashley and Buff 1986; Quinn 1988; Bart et al. 1994).

CONSERVATION ACCOMPLISHMENTS

South Carolina Stream Assessment (2006-2011) data have facilitated the calculation of standardized abundance (density) estimates for this species at multiple spatial strata including statewide, river basin, level-IV ecoregion, and “ecobasin” (ecoregion x river basin). These estimates, for the first time, provide an objective measure of current population status that will serve as a baseline for following future population trends and gauging the effectiveness of conservation actions.

Educational materials have been developed in order to raise public awareness of nongame species and their ecological importance to the natural history of South Carolina’s aquatic habitats, including:

- The Reel Art program creates a topic for secondary school students and judges the artists’ submissions (e.g. a list of the Piedmont Fishes of SC to select from as subjects for drawing or painting).
- We compiled information and photographs for the development of nongame fish description web pages which are currently in development.
- We developed the Blackwater River Guide and interactive Powerpoint.
 - <http://www.dnr.sc.gov/education/pdf/BlackwaterInteractivePoster.pdf>
 - <http://www.dnr.sc.gov/education/pdf/BlackwaterRivEdGuide.pdf>
- We developed and printed the Fish Species of Concern Coloring Book (2009).
 - <http://www.dnr.sc.gov/aquaticed/pdf/SCFishesofConcernColoringBook.pdf>

CONSERVATION RECOMMENDATIONS

- Use South Carolina Stream Assessment decision-support GIS modeling tools to identify levels and spatial distributions of critical habitat factors to sustain the species in geographic areas of interest.
- Use South Carolina Stream Assessment decision-support GIS modeling tools to identify priority regions and watersheds at greatest risk of decline in stream integrity.
- Describe life history and habitat requirements of the Snail Bullhead.
- Protect critical habitats from future development and further habitat degradation by following Best Management Practices (BMPs) and protecting and purchasing riparian areas.
- Promote land stewardship practices through educational programs both within critical habitats with healthy populations and other areas that contain available habitat.
- Encourage responsible land use planning.
- Consider this species' needs when participating in the environmental permit review process.
- Develop a Non-Game Fishes of South Carolina poster and other educational materials in order to raise public awareness of nongame species and their ecological importance to the natural history of South Carolina's aquatic habitats.
- Prevent the spread of Flathead Catfish and other non-indigenous aquatic species.
- Educate the public as to the discontents of stocking nonnative species.
- Continue to develop educational materials in order to raise public awareness of nongame species and their ecological importance to the natural history of South Carolina's aquatic habitats.
- Educate motor vehicle operators of the negative effects of crossing streams at multiple locations and using stream bottoms as trails.

MEASURES OF SUCCESS

Determining the distribution, life history, habitat needs, and Southeastern population structure and trends would represent a measure of success for this species. Methods that protect water quality are also likely to protect this species. In the event that more protective BMPs are implemented, population studies of this fish could assist in determining the effectiveness of those measures.

LITERATURE CITED

- Ashley, K.W. and B. Buff. 1986. Determination of current food habits of flathead catfish in the Cape Fear River. Final Report Submitted to the North Carolina Wildlife Resources Commission, Division of Boating and Inland Fisheries. Raleigh, North Carolina. 19 pp.
- Bart, H.L., M.S. Taylor, J.T. Harbaugh, J.W. Evans, S.L. Schleiger and W. Clark. 1994. New distribution records of Gulf Slope drainage fishes in the Ocmulgee River system, Georgia. Southeastern Fishes Council Proceedings. 30:4-9.

- Guire, C.R., L.E. Nichols and R.T. Rachels. 1984. Biological investigations of flathead catfish in the Cape Fear River. Proceedings of the Southeastern Association of Fish and Wildlife Agencies. 35(1981):607-621.
- Jelks, H.L., S.J. Walsh, N. M. Burkhead, S. Contreras-Balderas, E. Díaz-Pardo, D.A. Hendrickson, J. Lyons, N.E. Mandrak, F. McCormick, J.S. Nelson, S.P. Platania, B.A. Porter, C.B. Renaud, J.J. Schmitter-Soto, E.B. Taylor, and M.L. Warren, Jr. 2008. Conservation status of imperiled North American freshwater and diadromous Fishes. Fisheries 33 (5):372-407.
- Jenkins, R.E., and N.M. Burkhead. 1994. Freshwater Fishes of Virginia. American Fisheries Society. Bethesda, Maryland. 1079 pp.
- NatureServe. 2004. NatureServe Explorer: An online encyclopedia of life [web application]. Version 4.0. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: September 1, 2004).
- NatureServe. 2013. NatureServe Explorer: An online encyclopedia of life [web application]. Version 7.1. NatureServe, Arlington, Virginia. Available <http://www.natureserve.org/explorer>. (Accessed: March 26, 2013).
- Page, L.M. and B.M. Burr. 1991. A field guide to freshwater fishes: North America north of Mexico. Houghton Mifflin Company. Boston, Massachusetts. 432 pp.
- Quinn, S.P. 1988. Stomach contents of flathead catfish in the Flint River, Georgia. Proceedings of the 41st Annual Conference of Southeastern Association of Fish and Wildlife Agencies. 41(1987):85-92.
- Rohde, F.C., R.G., Arndt, D.G. Lindquist and J.F. Parnell. 1994. Freshwater Fishes of the Carolinas, Virginia, Maryland, and Delaware. The University of North Carolina Press. Chapel Hill, North Carolina. 222 pp.
- Warren, M.L., Jr., B.M. Burr, S.J. Walsh, H.L. Bart, Jr., R.C. Cashner, D.A. Etnier, B.J. Freeman, B.R. Kuhajda, R.L. Mayden, H.W. Robison, S.T. Ross and W.C. Starnes. 2000. Diversity, distribution, and conservation status of the native freshwater fishes of the southern United States. Fisheries 25(10):7-31.