Table 1. Summary of wood stork (*Mycteria americana*) nest monitoring data collected by South Carolina Department of Natural Resources staff, US Fish & Wildlife Service staff, and volunteers during 2011 - 2017.

Year	2011	2012	2013	2014	2015	2016	2017	All Years
Number of Monitored Nest Sites	81	311	427	396	412	322	232	2181
Average Fledglings per Nest Site	1.6	1.1	1.4	2.2	2.3	1.7	1.7	2.0
Average Fledgings per Successful Nest Site	2.1	1.9	2.0	2.6	2.5	2.3	2.0	2.2
0 Fledglings	18	122	144	49	48	88	52	521
1 Fledgling	11	53	55	28	26	42	23	238
2 Fledglings	38	107	163	126	144	98	100	776
3 Fledglings	14	28	59	163	168	73	51	556
4 Fledglings	0	1	6	30	29	21	6	93
			<u> </u>	<u> </u>	<u> </u>			<u> </u>
% Successful*	77%	61%	66%	88%	89%	74%	79%	77%

^{*} Number of successful nest sites divided by the total number of nest sites that were monitored. Successful is defined as producing at least one fledgling. A chick was considered to be a fledgling if it survived to at least 7 weeks of age.

Note: This data was collected as part of an ongoing monitoring project by the South Carolina Department of Natural Resources (SCDNR). Data is preliminary and may be revised. Please contact Christy Hand at SCDNR prior to using this data in reports or publications.

Figure 1. Number of wood stork ($Mycteria\ americana$) nests counted in South Carolina during annual censuses from 1981-2017.




Figure 2. Number of colonies in which wood storks ($Mycteria\ americana$) nested in South Carolina from 1981-2017

